



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

It is a beautiful sight to watch the Herons, some setting in the nests, some flying and some standing tall and majestic upon the dead snags, and it makes a picture long to be remembered with pleasure.

---

## THE LARK FINCH.

*Chondestes Grammica.*

BY J. A. SINGLEY, GIDDINGS, TEXAS.

The Lark Sparrow is a common summer resident here, arriving from the 1st to the 10th of April, and departing early in November. It can be found anywhere in the uplands and prairies during its stay with us ; but it does not frequent the bottoms. It is one of our best singers and occupies the same position here that the Grass Finch does in the northern states. The birds mate shortly after their arrival, and by May 1st nest building is in order, and eggs can be taken from the 10th of May until the last of July. The first nests are usually placed on the ground, but with the advance of the season as many or more nests will be found in trees and bushes than on the ground.

The nest varies considerably, sometimes being of grass throughout (this is the case when built on the ground) and again, of a mixture of grass and weeds lined with horse-hair. Very often an old Mockingbird nest is appropriated and reduced to smaller dimensions by filling up with grass and lining with hair. On one occasion I found where the Lark Finch had fitted up a nest of the Orchard Oriole, enlarging it to suit and depositing its eggs in the swinging nest.

Like most of the Sparrows, this species is sober colored. The upper parts grayish-brown, the middle of the back streaked with black : under parts white, tinged with brownish, a black blotch on the breast, top of head chestnut, with a median stripe of whitish. The lateral tail feathers are black, tipped with white. When flying, the tail is spread and the white tips show plainly, making it one of the easiest of the Sparrows to identify. The usual number of eggs in a set is four or five, but I have found as many as eight eggs in a set. Color, grayish-white, spotted and marked with dark, almost black lines and scrawls. The eggs resemble those of the Baltimore Oriole. Average, .85x.65.